

ILLUSTRATED REVIEW

Heparin-induced thrombocytopenia: An illustrated review

Jori May MD¹ | Brian Westbrook MD¹ | Adam Cuker MD, MS^{2,3} 

¹Department of Medicine, University of Alabama at Birmingham, Birmingham, Alabama, USA

²Department of Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

³Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

Correspondence

Jori May, Division of Hematology/Oncology, Department of Medicine, University of Alabama at Birmingham, 1720 2nd Ave South, NP 2503, Birmingham, AL 35294, USA.

Email: jemay@uabmc.edu

Handling Editor: Dr Michelle Sholzberg

Abstract

Heparin-induced thrombocytopenia (HIT) is an immune-mediated adverse drug effect from unfractionated or low-molecular-weight heparin that results in thrombocytopenia and potentially catastrophic thrombosis. HIT occurs due to the development of platelet-activating antibodies against multimolecular complexes of platelet factor 4 and heparin. Given the frequency of thrombocytopenia and heparin use among hospitalized patients, calculation of the 4Ts Score is recommended to identify patients at increased likelihood of HIT and direct further evaluation. In patients with an intermediate or high probability 4Ts Score, an immunoassay and functional assay are recommended to confirm or refute the diagnosis of HIT. Heparin avoidance and initiation of nonheparin anticoagulation are the mainstays of acute HIT management. In this illustrated review, we provide visual summaries of the diagnosis and management of HIT, highlighting connections between pathophysiology and clinical care as well as summarizing efforts in quality improvement in the field. We further emphasize common pitfalls and pearls in diagnosis and management to encourage evidence-based care. We include graphical representation of the unique challenges of HIT with cardiopulmonary bypass and also delineate autoimmune HIT and its subtypes.

KEYWORDS

anticoagulants, drug-related side effects and adverse reactions, heparin, quality improvement, thrombosis

Essentials

- Heparin-induced thrombocytopenia (HIT) is an immune adverse drug effect that results in low platelets and risk of thrombosis.
- Use of the 4Ts Score is recommended to assess patient risk of HIT.
- Sequential specialized testing is required to diagnose HIT.
- Heparin avoidance and use of nonheparin anticoagulation are the mainstays of HIT management.

Heparin-induced thrombocytopenia (HIT)

Definition: An immune-mediated adverse drug event to heparin that results in **thrombocytopenia** and potentially **catastrophic thrombosis**.

This illustrated review will discuss the **diagnosis and management** of HIT, highlighting connections between  **Pathophysiology** and clinical care, and emphasizing central  **Pitfalls** and  **Pearls** to encourage evidence-based care.

Incidence

↑ incidence with **major surgery**, particularly cardiac²


1 in 1500 hospitalizations¹
in the United States

↑ incidence with **unfractionated heparin** (3%) vs. low molecular weight (<1%)^{3,4}

↓ incidence with **shorter duration of exposure** to heparin, specifically <5 days⁵

Clinical manifestations & outcomes¹

Compared to hospital discharges without HIT



↓ *Single institution reports*⁹

ratio
Venous:Arterial
1:1 to 4:1

3-8%

6-50%

Rare manifestations



Disseminated intravascular coagulation (DIC)¹⁰



Injection site necrotizing skin lesions¹¹




Anaphylactoid reactions to IV heparin bolus¹¹


aOR, adjusted odds ratio; CI, confidence interval

Pathophysiology


The Key Players



How does the anti-PF4/heparin antibody cause HIT?



Why don't all anti-PF4/heparin antibodies cause HIT?



Diagnosis, Step 1: The 4Ts Score


4Ts Score²¹ = Pre-test scoring system to assess the probability of HIT

Calculated as a sum of points from 4 components:

① Thrombocytopenia

Points:

- 2** — Platelet fall > 50% *and* platelet nadir $\geq 20 \times 10^9/L$
- 1** — Platelet fall 30%-50% *or* Platelet nadir $10-19 \times 10^9/L$
- 0** — Platelet fall < 30% *or* Platelet nadir < $10 \times 10^9/L$



! Calculate using *peak* platelet count *after* heparin started (i.e., not necessarily the day heparin started)

! % decline is used, not an absolute platelet threshold, as HIT can occur without thrombocytopenia if the peak count is high (e.g., after surgery, myeloproliferative neoplasm).²¹


② Timing of platelet count fall

Points:

- 2** — Fall starts days 5-10 (timing is clear) *or* Fall starts ≤ 1 day + heparin exposure within 30 days
- 1** — Fall starts days 5-10 (timing is unclear) *or* Fall starts after day 10 *or* Fall starts ≤ 1 day + heparin exposure 30-100 days ago
- 0** — Fall starts ≤ 4 days without recent heparin exposure

Visualization of points:

Line color correlates with # points listed above



! Calculated based on the *first day* of consistent platelet fall.

! In many patients, platelet fall can be *biphasic*, with an initial decline due to a non-HIT insult (e.g. surgery), followed by a rebound and a second, HIT-induced decline.

! It takes 5-10 days to produce new IgG antibodies. Immediate HIT onset can occur in those with recent exposure due to presence of previously formed antibodies.^{23,24}

④ Other causes of thrombocytopenia

Points:

- 2** None apparent
- 1** Possible
- 0** Definite

! This variable can be challenging to determine and leads to variation between providers in score determination.²⁵

! **HEP score:**²⁶ An alternate score with similar criteria but better performance with less experienced clinicians, likely due to inclusion of defined "other causes of thrombocytopenia."²⁷

1. Chronic thrombocytopenia
2. Newly initiated non-heparin med known to cause thrombocytopenia
3. Severe infection
4. Severe disseminated intravascular coagulation (DIC)
5. Indwelling intra-arterial device
6. Cardiopulmonary bypass within 96 hours

Total 4Ts Score
= probability of HIT²²

- ≥ 6 High (64%)**
- 4-5 Intermediate (14%)**
- ≤ 3 Low (0.2%)**

③ Thrombosis or other sequelae

Points:

- 2** New thrombosis; skin necrosis; acute systemic reaction post-intravenous UFH bolus
- 1** Progressive or recurrent thrombosis; non-necrotizing (erythematous) skin lesions at heparin injection sites; suspected thrombosis (not proven)
- 0** None

! Includes thrombosis that occurs or progresses in the period of platelet count decline *after* heparin exposure (i.e., not thrombosis that occurred before heparin exposure)

Diagnosis, Step 2a: Laboratory diagnosis, *Immunoassays*

The laboratory diagnosis of HIT is classically divided into 2 steps:

(a) an **immunoassay** and (b) a **functional assay**.


Enzyme Immunoassay (EIA)

The EIA is the prototypical immunoassay, and is used as the standard immunoassay throughout this manuscript

Purpose: Determine if anti-PF4/heparin antibodies are *present*, and if so, their abundance

Performance: High sensitivity (0.97, 95% CI 0.95-0.99), limited specificity (0.82, 95% CI 0.90-0.84)²⁸

 Specificity varies based on assay used, increasing with IgG-specific assays^{28,29} and higher OD thresholds.³⁰



 Higher OD correlates with higher HIT likelihood³⁰ and thrombotic risk.³¹

How it works: (A) The plate well is coated with multimolecular complexes of PF4/heparin (or PF4/heparin-like polyanions). Patient serum or plasma is added and anti-PF4/heparin antibodies bind to the complex. (B) Tagged anti-human antibody is added and binds anti-PF4/heparin antibodies. A substrate is added to elicit color change. (C) Degree of color change (optical density, OD) increases with increasing concentration of anti-PF4/heparin antibodies.

If EIA is positive, proceed with functional assay (see next page, "Laboratory diagnosis, Functional assays").

If EIA is negative, HIT is essentially excluded.

Additional immunoassay options: Rapid Immunoassays

Multiple types of immunoassays exist, all with different performance characteristics and different international availability. Selected examples of rapid immunoassays are included below.

Purpose: Similar to EIA but provide results in ≤ 30 minutes (EIA is often batched, takes hours to perform)

Performance: Varies by assay

		Sensitivity (95% CI)	Specificity (95% CI)	How it works: Method of anti-PF4/heparin detection	Ab class detected
Particle gel immunoassay ³²	PaGIA	0.98 (0.94-1.00)	0.88 (0.91-0.96)	 Polymers coated with PF4/heparin complexes	IgG1
IgG-specific chemiluminescence assay ³²	IgG CIA	0.95 (0.90-1.00)	0.94 (0.89-0.99)	 Magnetic particles coated with PF4/polyvinyl sulfonate	IgG
Lateral flow immunoassay ³²	LFIA	0.97 (0.92-1.00)	0.91 (0.86-0.96)	 PF4-polyanion complexes	IgG
Latex immunoturbidimetric assay ³³	LIA	0.974	0.94	 Competitive inhibition of nanoparticles coated with HIT-like antibodies	IgG, IgA, IgM

There is limited guidance on how rapid immunoassays should be incorporated into diagnostic algorithms, and whether additional confirmatory laboratory testing is needed.³²⁻³³

Diagnosis, Step 2b: Laboratory diagnosis, *Functional assays*


¹⁴C-serotonin release assay (SRA)

The SRA is the "gold standard" functional assay^{34,35} and is used as the standard functional assay throughout this manuscript

Purpose: Determine if anti-PF4/heparin antibodies are *pathogenic* (i.e. activate platelets)

Performance: High sensitivity (~0.95), high specificity (~0.95)³⁵

 Due to variation in assay performance and reporting practices,³⁶ real-world performance may vary from literature reports.




How it works:^{34,35} (A) Patient serum is combined with washed donor platelets radiolabeled with ¹⁴C-serotonin. (B) At least two different doses of heparin are added. At therapeutic dose, the HIT antigen forms and is recognized by the anti-PF4/heparin antibody. At high dose, heparin disrupts the multimolecular HIT antigen^{34,39} so the anti-PF4/heparin antibody cannot bind. (C) When the antigen/antibody complex binds to the platelet Fc receptor, platelets activate and release granules containing ¹⁴C-serotonin. Radioactivity is measured and converted to a % ¹⁴C-serotonin release. (D) A diagnosis of HIT (i.e. positive SRA) is generally characterized by strong activation (high % serotonin release) with therapeutic dose heparin *plus* inhibited activation (low % serotonin release) with high dose heparin. A negative SRA shows no activation at either concentration. An indeterminate SRA shows activation at both heparin concentrations.

Additional functional assays

Multiple functional assays exist that differ by the method used for measuring platelet activation:


Heparin-induced platelet activation assay (HIPA)⁴⁰

Measurement of activation:
Visual assessment of decreased turbidity with platelet activation



P-selectin expression assay (PEA)⁴¹⁻⁴³

Measurement of activation:
Platelet P-selectin (CD62p) expression by flow cytometry



Diagnosis: Putting it all together



Why use risk scores before laboratory testing?

- Thrombocytopenia is common in hospitalized patients, and it is rarely due to HIT⁴⁴
- The 4Ts Score has excellent *negative* predictive value (can exclude HIT)²²
- EIA is sensitive, not specific. False positives are common.



Choosing Wisely recommendation to avoid laboratory testing in patients with low probability 4Ts Score^{45,46}



Why not wait for the laboratory results to change anticoagulation?

Without appropriate therapy in a patient with HIT, the daily risk of complications (combined risk of thromboembolism, amputation, and death) is 6.1%.⁴⁷



Risk of HIT can be further stratified based on EIA optical density (OD) and 4Ts Score:

EIA OD	4Ts Score		without EIA OD
	4-5	6-8	
<0.6	0%	0%	0%
0.6-1.49	5-25%	16%	68%
1.5-1.99	50%	53%	93%
2+	90%	92%	99%

without 4Ts Score


Lighter shades indicate probability of the individual component (4Ts Score without EIA OD,²² EIA OD without 4Ts Score⁴⁸). Darker shades indicate probability accounting for combination of 4Ts Score and EIA OD.^{49,50} Exact probabilities will vary with different EIA assays.



In patients with a high probability 4Ts score and a strongly positive EIA, a functional assay may not be necessary.⁵¹




Rarely, functional assays are falsely negative.⁵² In patients with a high 4Ts score and high EIA OD, treatment for HIT can be considered despite a negative functional assay.⁵¹



HIT phases & management

The phases of HIT: The timeline of HIT diagnosis and progression can be defined in phases.⁵³ Phases are defined by 4 clinical parameters: (1) platelet count, (2) functional assay result, (3) EIA result, and (4) risk of thrombosis. Red indicates the feature is abnormal, yellow indicates the feature is unclear (e.g., in suspected HIT, EIA result is unknown), and green indicates the feature has returned to baseline.



Who requires platelet monitoring for HIT?

Recommendations⁵¹ based on risk influenced by (1) type of heparin product and (2) patient population

Estimated risk of HIT	Platelet count monitoring*	Schedule assumes no heparin exposure within 30 days. If monitoring, includes day 4-14 after heparin exposure (day 0). Consider a platelet count before day 4 to identify baseline in case of decline.
 Low <0.1%	None	
 Intermediate 0.1-1%	Consider every 2-3 days	
 High >1%	Consider at least every other day	

	LMWH	UFH*	Fondaparinux
Medical			
Obstetric			
Minor surgery			
Major surgery			
Minor trauma			
Major trauma			

*If exposed to LMWH and UFH, follow guidance for UFH


2 key principles of HIT management

① Avoid heparin

Duration of heparin avoidance: Indefinite avoidance is recommended. Brief exposure can be considered in Subacute HIT B and Remote HIT in specific scenarios (See *HIT in cardiopulmonary bypass*).

② Use non-heparin anticoagulation

Duration of non-heparin anticoagulation: Depends on the presence or absence of thrombosis.⁵¹ For details of agent selection, see *Selection of non-heparin anticoagulation & other therapies*



Selection of non-heparin anticoagulation & other therapies

Anticoagulation selection in **Acute HIT**

Dependent on 4 clinical criteria

		1) Clinically unstable	2) Life- or limb-threatening ischemia	3) Renal dysfunction CrCl <30ml/min	4) Liver dysfunction Bili >1.5mg/dL
Oral	Oral Xa inhibitors*	●	●	●	●
	Vitamin K antagonist (VKA)	●	●	●	●
IV	Argatroban	●	●	●	●
	Bivalirudin	●	●	● [^]	●
SQ	Danaparoid	●	●	●	●
	Fondaparinux ^o	●	●	● [~]	●

- Agent is preferred
- Agent is not preferred, but can be considered based on availability, risk/benefit
- Agent is not recommended

 Laboratory monitoring required

* Existing data with rivaroxaban, apixaban

[^] If argatroban not available, can use with close monitoring due to accumulation risk

^o Trivial risk of reported HIT,⁵⁴ but has been demonstrated to be safe for use in acute HIT^{55,56}


[~] Use in renal dysfunction has been reported^{57,58}

^o Trivial risk of reported HIT,⁵⁴ but has been demonstrated to be safe for use in acute HIT^{55,56}

[~] Use in renal dysfunction has been reported^{57,58}

Bili, bilirubin; CrCl, creatinine clearance; IV, intravenous; SQ, subcutaneous


Oral anticoagulation pearls^{51,59}



Additional treatment considerations

Intravenous immunoglobulin (IVIg)

- Hypothesized to competitively inhibit binding of anti-PF4/heparin antibody to platelet FcR.⁶²
- Can be considered in cases of **severe and/or refractory thrombosis or thrombocytopenia**.



Anti-PF4/heparin antibody complex

IVIg

Thrombosis

Transfusion


STOP Platelet transfusion?

- Routine use **NOT** recommended due to concern it may "fuel the fire" and increase thrombotic risk, particularly arterial
- Can be considered in patients with active bleeding or prior to urgent procedures.^{51,5}


HIT in cardiopulmonary bypass (CPB)

CPB is associated with a higher risk of HIT than other surgeries because (1) it requires high doses of anticoagulation, most commonly unfractionated heparin, and (2) it causes intense platelet activation and PF4 release.


Incidence & diagnosis



Platelet count trajectory in diagnosis



Performing CPB in patients with a history of HIT^{51,65}





Autoimmune HIT (aHIT)

What is autoimmune HIT?


- Rarely, the clinical manifestations of HIT can develop via **heparin-independent platelet activation**.
- Autoimmune HIT (aHIT) refers to any HIT-related platelet count fall that begins or persists in the absence of heparin.^{70,71}
- Six subtypes of aHIT have been described.
- Notably, spontaneous HIT has been reported to occur most commonly after knee replacement or infection.⁷¹

Pathophysiology




(1) Positively charged PF4 tetramers repel one another. (2) In aHIT, a non-heparin molecule overcomes this electrostatic repulsion, bringing individual PF4 tetramers together to form large multimolecular PF4 complexes. The exact non-heparin molecule able to perform this action is debated and may vary based on aHIT subtype, but proposed examples include HIT "super-antibodies" that bridge two PF4 tetramers and/or highly anionic molecules which could include chondroitin sulfate (released from joint cartilage in joint surgery), polyphosphate, and/or nucleotides such as DNA and RNA.^{70,71} (3) The PF4 complexes then serve as antigens bound by anti-PF4/polyanion IgG, which (4) trigger activation of platelets and other cell types.

Principles of diagnosis & management of aHIT syndromes are ***the same as classical HIT***, with some unique considerations:



Quality improvement in diagnosis & management

The diagnosis and management of HIT are ripe for opportunities to improve the delivery of evidence-based, cost-effective care. Below are examples of successful initiatives, with discussion of the details of the intervention and its impact.

	Goal	Intervention	Impact
Heparin exposure	 Decrease exposure to unfractionated heparin (UFH) to decrease incidence of HIT	 The Avoid Heparin Initiative:⁷² Replacement of UFH with LMWH for prophylactic and therapeutic indications	↓ total HIT incidence by 79% (10.7 to 2.2 per 10,000 admission) ↓ HITT incidence by 91% (4.6 to 0.4 cases per 10,000 admissions) ↓ HIT-related costs by \$266,938 per year
Suspected HIT	 Decrease unnecessary testing for HIT	 Clinical decision support requiring 4Ts Score calculation with all HIT assay^{73,74}	↓ proportion of tested patients with low 4Ts scores from 66% to 56%⁷³ ↓ screening and functional assay testing by 37.5% and 85%, respectively⁷⁴
Acute HIT	 Increase appropriate use of non-heparin anticoagulants	 Creation of an anticoagulation stewardship program,⁷⁵ with a focus on anticoagulation in HIT	↓ median duration of IV direct thrombin inhibitor use from 9.9 days to 7.3 days ↓ time from diagnosis to discharge in acute HIT from 12.8 days to 9.2 days ↑ fondaparinux and oral Xa inhibitor prescribing rates in acute HIT from 40.7% to 62.2%

ACKNOWLEDGMENTS

The authors thank Douglas Cines, MD, for his expert review of the manuscript. Illustrations created with BioRender.com.

FUNDING

The authors received no funding for this study.

AUTHOR CONTRIBUTIONS

J.M. conceptualized the manuscript and created the initial content and visualization. B.W. and A.C. provided critical review and revision.

RELATIONSHIP DISCLOSURE

J.M. and B.W. report no relevant conflicts of interest. A.C. has served as a consultant for Synergy, MingSight, and the New York Blood Center and has received authorship royalties from UpToDate.

TWITTER

Adam Cuker  @CukerMd

REFERENCES

- [1] Dhakal B, Kreuziger LB, Rein L, Kleman A, Fraser R, Aster RH, et al. Disease burden, complication rates, and health-care costs of heparin-induced thrombocytopenia in the USA: a population-based study. *Lancet Haematol*. 2018;5:e220–31.
- [2] Pishko AM, Cuker A. Heparin-induced thrombocytopenia in cardiac surgery patients. *Semin Thromb Hemost*. 2017;43:691–8.
- [3] Warkentin TE, Levine MN, Hirsh J, Horsewood P, Roberts RS, Gent M, et al. Heparin-induced thrombocytopenia in patients treated with low-molecular-weight heparin or unfractionated heparin. *N Engl J Med*. 1995;332:1330–5.
- [4] Martel N, Lee J, Wells PS. Risk for heparin-induced thrombocytopenia with unfractionated and low-molecular-weight heparin thromboprophylaxis: a meta-analysis. *Blood*. 2005;106:2710–5.
- [5] Smythe MA, Koerber JM, Mattson JC. The incidence of recognized heparin-induced thrombocytopenia in a large, tertiary care teaching hospital. *Chest*. 2007;131:1644–9.
- [6] Warkentin TE, Sheppard J-AI, Whitlock RP. Temporal presentations of heparin-induced thrombocytopenia following cardiac surgery: a single-center, retrospective cohort study. *J Thromb Haemost*. 2022;20:2601–16.
- [7] Warkentin TE, Sheppard J-AI, Heels-Ansdell D, Marshall JC, McIntyre L, Rocha MG, et al. Heparin-induced thrombocytopenia in medical surgical critical illness. *Chest*. 2013;144:848–58.
- [8] Gruel Y, Vayne C, Rollin J, Weber P, Faille D, Bauters A, et al. Comparative analysis of a French prospective series of 144 patients with heparin-induced thrombocytopenia (FRIGTIH) and the literature. *Thromb Haemost*. 2020;120:1096–107.
- [9] Arepally GM, Padmanabhan A. Heparin-induced thrombocytopenia: a focus on thrombosis. *Arterioscler Thromb Vasc Biol*. 2021;41:141–52.
- [10] Warkentin TE. Heparin-induced thrombocytopenia. *Curr Opin Crit Care*. 2015;21:576–85.
- [11] Warkentin TE, Roberts RS, Hirsh J, Kelton JG. Heparin-induced skin lesions and other unusual sequelae of the heparin-induced thrombocytopenia syndrome: a nested cohort study. *Chest*. 2005;127:1857–61.
- [12] Bock PE, Luscombe M, Marshall SE, Pepper DS, Holbrook JJ. The multiple complexes formed by the interaction of platelet factor 4 with heparin. *Biochem J*. 1980;191:769–76.
- [13] Chong BH, Fawaz I, Chesterman CN, Berndt MC. Heparin-induced thrombocytopenia: mechanism of interaction of the heparin-dependent antibody with platelets. *Br J Haematol*. 1989;73:235–40.
- [14] Gollomp K, Kim M, Johnston I, Hayes V, Welsh J, Arepally GM, et al. Neutrophil accumulation and NET release contribute to thrombosis in HIT. *JCI Insight*. 2018;3:e99445.
- [15] Khandelwal S, Barnes A, Rauova L, Sarkar A, Rux AH, Yarovi SV, et al. Complement mediates binding and procoagulant effects of ultralarge HIT immune complexes. *Blood*. 2021;138:2106–16.
- [16] Suh JS, Malik MI, Aster RH, Visentin GP. Characterization of the humoral immune response in heparin-induced thrombocytopenia. *Am J Hematol*. 1997;54:196–201.
- [17] Lindhoff-Last E, Gerdson F, Ackermann H, Bauersachs R. Determination of heparin-platelet factor 4-IgG antibodies improves diagnosis of heparin-induced thrombocytopenia. *Br J Haematol*. 2001;113:886–90.
- [18] Greinacher A, Alban S, Omer-Adam MA, Weitschies W, Warkentin TE. Heparin-induced thrombocytopenia: a stoichiometry-based model to explain the differing immunogenicities of unfractionated heparin, low-molecular-weight heparin, and fondaparinux in different clinical settings. *Thromb Res*. 2008;122:211–20.
- [19] Cines DB, Rauova L, Arepally G, Reilly MP, McKenzie SE, Sachais BS, et al. Heparin-induced thrombocytopenia: an autoimmune disorder regulated through dynamic autoantigen assembly/disassembly. *J Clin Apher*. 2007;22:31–6.
- [20] Suvarna S, Espinasse B, Qi R, Lubica R, Poncz M, Cines DB, et al. Determinants of PF4/heparin immunogenicity. *Blood*. 2007;110:4253–60.
- [21] Lo GK, Juhl D, Warkentin TE, Sigouin CS, Eichler P, Greinacher A. Evaluation of pretest clinical score (4 Ts) for the diagnosis of heparin-induced thrombocytopenia in two clinical settings. *J Thromb Haemost*. 2006;4:759–65.
- [22] Cuker A, Gimotty PA, Crowther MA, Warkentin TE. Predictive value of the 4Ts scoring system for heparin-induced thrombocytopenia: a systematic review and meta-analysis. *Blood*. 2012;120:4160–7.
- [23] Warkentin TE, Kelton JG. Temporal aspects of heparin-induced thrombocytopenia. *N Engl J Med*. 2001;344:1286–92.
- [24] Lefler DS, Cuker A, Linkins LA, Warkentin TE, Pishko AM. Maximum 24-hour platelet count fall: metric for improving the diagnosis of heparin-induced thrombocytopenia among patients with intermediate probability 4Ts scores. *J Thromb Haemost*. 2020;18:2018–24.
- [25] Crowther M, Cook D, Guyatt G, Zytaruk N, McDonald E, Williamson D, et al. Heparin-induced thrombocytopenia in the critically ill: interpreting the 4Ts test in a randomized trial. *J Crit Care*. 2014;29:470.e7, 15.
- [26] Cuker A, Arepally G, Crowther MA, Rice L, Datko F, Hook K, et al. The HIT Expert Probability (HEP) Score: a novel pre-test probability model for heparin-induced thrombocytopenia based on broad expert opinion. *J Thromb Haemost*. 2010;8:2642–50.
- [27] Pishko AM, Fardin S, Lefler DS, Paydary K, Vega R, Arepally GM, et al. Prospective comparison of the HEP score and 4Ts score for the diagnosis of heparin-induced thrombocytopenia. *Blood Adv*. 2018;2:3155–62.
- [28] Hussein-zadeh HD, Gimotty PA, Pishko AM, Buckley M, Warkentin TE, Cuker A. Diagnostic accuracy of IgG-specific versus polyspecific enzyme-linked immunoassays in heparin-induced thrombocytopenia: a systematic review and meta-analysis. *J Thromb Haemost*. 2017;15:1203–12.
- [29] Cuker A, Ortel TL. ASH evidence-based guidelines: is the IgG-specific anti-PF4/heparin ELISA superior to the polyspecific ELISA in the laboratory diagnosis of HIT? *Hematology Am Soc Hematol Educ Program*. 2009;250, 2.
- [30] Warkentin TE, Sheppard JI, Moore JC, Sigouin CS, Kelton JG. Quantitative interpretation of optical density measurements using

- PF4-dependent enzyme-immunoassays. *J Thromb Haemost.* 2008;6:1304–12.
- [31] Zwicker JL, Uhl L, Huang WY, Shaz BH, Bauer KA. Thrombosis and ELISA optical density values in hospitalized patients with heparin-induced thrombocytopenia. *J Thromb Haemost.* 2004;2:2133–7.
 - [32] Sun L, Gimotty PA, Lakshmanan S, Cuker A. Diagnostic accuracy of rapid immunoassays for heparin-induced thrombocytopenia. A systematic review and meta-analysis. *Thromb Haemost.* 2016;115:1044–55.
 - [33] Warkentin TE, Sheppard J-AI, Linkins L-A, Arnold DM, Nazy I. Performance characteristics of an automated latex immunoturbidimetric assay [HemosIL®HIT-Ab_(PF4-H)] for the diagnosis of immune heparin-induced thrombocytopenia. *Thromb Res.* 2017;153:108–17.
 - [34] Sheridan D, Carter C, Kelton JG. A diagnostic test for heparin-induced thrombocytopenia. *Blood.* 1986;67:27–30.
 - [35] Warkentin TE, Arnold DM, Nazy I, Kelton JG. The platelet serotonin-release assay. *Am J Hematol.* 2015;90:564–72.
 - [36] Price EA, Hayward CP, Moffat KA, Moore JC, Warkentin TE, Zehnder JL. Laboratory testing for heparin-induced thrombocytopenia is inconsistent in North America: a survey of North American specialized coagulation laboratories. *Thromb Haemost.* 2007;98:1357–61.
 - [37] Warkentin TE, Hayward CP, Smith CA, Kelly PM, Kelton JG. Determinants of donor platelet variability when testing for heparin-induced thrombocytopenia. *J Lab Clin Med.* 1992;120:371–9.
 - [38] Pötzsch B, Keller M, Madlener K, Müller-Berghaus G. The use of heparinase improves the specificity of crossreactivity testing in heparin-induced thrombocytopenia. *Thromb Haemost.* 1996;76:1121.
 - [39] Greinacher A, Alban S, Dummel V, Franz G, Mueller-Eckhardt C. Characterization of the structural requirements for a carbohydrate based anticoagulant with a reduced risk of inducing the immunological type of heparin-associated thrombocytopenia. *Thromb Haemost.* 1995;74:886–92.
 - [40] Greinacher A, Michels I, Kiefel V, Mueller-Eckhardt C. A rapid and sensitive test for diagnosing heparin-associated thrombocytopenia. *Thromb Haemost.* 1991;66:734–6.
 - [41] Padmanabhan A, Jones CG, Curtis BR, Bougie DW, Sullivan MJ, Peswani N, et al. A novel PF4-dependent platelet activation assay identifies patients likely to have heparin-induced thrombocytopenia/thrombosis. *Chest.* 2016;150:506–15.
 - [42] Padmanabhan A, Jones CG, Bougie DW, Curtis BR, McFarland JG, Wang D, et al. A modified PF4-dependent, CD62p expression assay selectively detects serotonin-releasing antibodies in patients suspected of HIT. *Thromb Haemost.* 2015;114:1322–3.
 - [43] Samuelson Bannow B, Warad DM, Jones CG, Pechauer SM, Curtis BR, Bougie DW, et al. A prospective, blinded study of a PF4-dependent assay for HIT diagnosis. *Blood.* 2021;137:1082–9.
 - [44] Crowther MA, Cook DJ, Meade MO, Griffith LE, Guyatt GH, Arnold DM, et al. Thrombocytopenia in medical-surgical critically ill patients: prevalence, incidence, and risk factors. *J Crit Care.* 2005;20:348–53.
 - [45] Li A, Hicks LK, Fan E. Things we do for no reason: HIT testing in low probability patients. *J Hosp Med.* 2019;14:374–81.
 - [46] American Society of Hematology. ten things physicians and patients should question. Choosing wisely: an initiative of the ABIM Foundation. <https://www.choosingwisely.org/societies/american-society-of-hematology/>; 2014. [accessed July 1, 2022].
 - [47] Greinacher A, Eichler P, Lubenow N, Kwasny H, Luz M. Heparin-induced thrombocytopenia with thromboembolic complications: meta-analysis of 2 prospective trials to assess the value of parenteral treatment with lepirudin and its therapeutic aPTT range. *Blood.* 2000;96:846–51.
 - [48] Warkentin TE. How I diagnose and manage HIT. *Hematology Am Soc Hematol Educ Program.* 2011;2011:143–9.
 - [49] Raschke RA, Gallo T, Curry SC, Whiting T, Padilla-Jones A, Warkentin TE, et al. Clinical effectiveness of a Bayesian algorithm for the diagnosis and management of heparin-induced thrombocytopenia. *J Thromb Haemost.* 2017;15:1640–5.
 - [50] Cuker A. Clinical and laboratory diagnosis of heparin-induced thrombocytopenia: an integrated approach. *Semin Thromb Hemost.* 2014;40:106–14.
 - [51] Cuker A, Arepally GM, Chong BH, Cines DB, Greinacher A, Gruel Y, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: heparin-induced thrombocytopenia. *Blood Adv.* 2018;2:3360–92.
 - [52] Warkentin TE, Nazy I, Sheppard J-AI, Smith JW, Kelton JG, Arnold DM. Serotonin-release assay-negative heparin-induced thrombocytopenia. *Am J Hematol.* 2020;95:38–47.
 - [53] Cuker A. Management of the multiple phases of heparin-induced thrombocytopenia. *Thromb Haemost.* 2016;116:835–42.
 - [54] Warkentin TE, Maurer BT, Aster RH. Heparin-induced thrombocytopenia associated with fondaparinux. *N Engl J Med.* 2007;356:2653–5.
 - [55] Kang M, Alahmadi M, Sawh S, Kovacs MJ, Lazo-Langner A. Fondaparinux for the treatment of suspected heparin-induced thrombocytopenia: a propensity score-matched study. *Blood.* 2015;125:924–9.
 - [56] Schindewolf M, Steindl J, Beyer-Westendorf J, Schellong S, Dohmen PM, Brachmann J, et al. Use of fondaparinux off-label or approved anticoagulants for management of heparin-induced thrombocytopenia. *J Am Coll Cardiol.* 2017;70:2636–48.
 - [57] Mahieu E, Claes K, Jacquemin M, Evenepoel P, Op De Beek K, Bogaert AM, et al. Anticoagulation with fondaparinux for hemodiafiltration in patients with heparin-induced thrombocytopenia: dose-finding study and safety evaluation. *Artif Organs.* 2013;37:482–7.
 - [58] Ghaziri D, Dehaini H, Msheik M, Bahmad M, Zorkot M, Saad GA. Novel fondaparinux protocol for anticoagulation therapy in adults with renal failure and suspected heparin-induced thrombocytopenia: a retrospective review of institutional protocol. *BMC Pharmacol Toxicol.* 2023;24:2.
 - [59] Linkins L-A, Dans AL, Moores LK, Bona R, Davidson BL, Schulman S, et al. Treatment and prevention of heparin-induced thrombocytopenia: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest.* 2012;141:e495S–530S.
 - [60] Warkentin TE, Elavathil LJ, Hayward CP, Johnston MA, Russett JI, Kelton JG. The pathogenesis of venous limb gangrene associated with heparin-induced thrombocytopenia. *Ann Intern Med.* 1997;127:804–12.
 - [61] Srinivasan AF, Rice L, Bartholomew JR, Rangaswamy C, La Perna L, Thompson JE, et al. Warfarin-induced skin necrosis and venous limb gangrene in the setting of heparin-induced thrombocytopenia. *Arch Intern Med.* 2004;164:66–70.
 - [62] Warkentin TE. High-dose intravenous immunoglobulin for the treatment and prevention of heparin-induced thrombocytopenia: a review. *Expert Rev Hematol.* 2019;12:685–98.
 - [63] Welsby IJ, Krakow EF, Heit JA, Williams EC, Arepally GM, Bar-Yosef S, et al. The association of anti-platelet factor 4/heparin antibodies with early and delayed thromboembolism after cardiac surgery. *J Thromb Haemost.* 2017;15:57–65.
 - [64] Warkentin TE, Sheppard JA, Horsewood P, Simpson PJ, Moore JC, Kelton JG. Impact of the patient population on the risk for heparin-induced thrombocytopenia. *Blood.* 2000;96:1703–8.
 - [65] Pishko AM, Cuker A. Heparin-induced thrombocytopenia and cardiovascular surgery. *Hematology Am Soc Hematol Educ Program.* 2021;2021:536–44.
 - [66] Lillo-Le Louët A, Boutouyrie P, Alhenc-Gelas M, Le Beller C, Gautier I, Aiach M, et al. Diagnostic score for heparin-induced thrombocytopenia after cardiopulmonary bypass. *J Thromb Haemost.* 2004;2:1882–8.

- [67] Pouplard C, May MA, Regina S, Marchand M, Fusiardi J, Gruel Y. Changes in platelet count after cardiac surgery can effectively predict the development of pathogenic heparin-dependent antibodies. *Br J Haematol*. 2005;128:837–41.
- [68] Warkentin TE. Acute intraoperative HIT during heart surgery: why so rare? *Thromb Res*. 2016;146:110–2.
- [69] Padmanabhan A, Connelly-Smith L, Aqui N, Balogun RA, Klingel R, Meyer E, et al. Guidelines on the use of therapeutic apheresis in clinical practice - evidence-based approach from the writing Committee of the American Society for Apheresis: the Eighth Special Issue. *J Clin Apher*. 2019;34:171–354.
- [70] Greinacher A, Selleng K, Warkentin TE. Autoimmune heparin-induced thrombocytopenia. *J Thromb Haemost*. 2017;15:2099–114.
- [71] Warkentin TE, Greinacher A. Spontaneous HIT syndrome: knee replacement, infection, and parallels with vaccine-induced immune thrombotic thrombocytopenia. *Thromb Res*. 2021;204:40–51.
- [72] McGowan KE, Makari J, Diamantouros A, Bucci C, Rempel P, Selby R, et al. Reducing the hospital burden of heparin-induced thrombocytopenia: impact of an avoid-heparin program. *Blood*. 2016;127:1954–9.
- [73] Samuelson BT, Glynn E, Holmes M, White AA, Martin DB, Garcia D. Use of a computer-based provider order entry (CPOE) intervention to optimize laboratory testing in patients with suspected heparin-induced thrombocytopenia. *Thromb Res*. 2015;136:928–31.
- [74] Lim MY, Foster J, Rourke A, Greenberg CS. Initial and long term impact of a multi-disciplinary task force in the diagnosis and management of heparin-induced thrombocytopenia. *J Thromb Thrombolysis*. 2018;45:130–4.
- [75] Lee J, Lindsley J, Chasler J, Streiff MB, Naik R, Shanbhag S, et al. The impact of an antithrombotic stewardship program on heparin-induced thrombocytopenia management. *J Pharm Pract*. 2022: 8971900221116185.